

Latest Bernese advances in Lunar geodesy

Stefano Bertone

Daniel Arnold Adrian Jäggi Valère Girardin Leos Mervart

Astronomical Institute, University of Bern, Switzerland

European Planetary Science Congress 2017
Riga, October 11, 2017

Outline

Lunar Geodesy

GRAIL @ AIUB

- Observables and modeling
- Orbit recovery
- Gravity field determination
- GRAIL+LLR

Conclusions



Lunar Geodesy




LLR

rotation, librations,
distance




LRO

topography, rotation




GRAIL

gravity, tides



Credits: NASA/GSFC/JPL/Colorado School of Mines/MIT

The GRAIL mission



GRAIL


(Gravity Recovery and Interior Laboratory)

M.T. Zuber et al, Science, 2013

March - December 2012

Polar orbit, altitude 25-50 km

- Understand the internal structure and history of the Moon
- Homogeneous coverage of the far side (GRACE heritage)




pre-GRAIL solutions:

LP165A (Lunar Prospector)

SGM150J (Selene)

The GRAIL mission



GRAIL


(Gravity Recovery and Interior Laboratory)

M.T. Zuber et al, Science, 2013

March - December 2012

Polar orbit, altitude 25-50 km

- Understand the internal structure and history of the Moon
- Homogeneous coverage of the far side (GRACE heritage)




Latest NASA solutions:

GL1500E (JPL)

GRGM1200A (GSFC)

The GRAIL mission: Satellite signals



- S-band two-way link (2.3 GHz)
- X-band one-way link (8.4 GHz)

One-way X-band Doppler (nominal 0.08 mm/s):

- emission/reception by different clocks
- smaller onboard antenna

Two-way S-band Doppler (nominal 0.3 mm/s):

- emission/reception by the same clock (DSN)
- larger onboard antenna


Ka-Band Range Rate (KBRR) (nominal $0.03\mu\text{m/s}$):

- Derived from Lunar Gravity Ranging System one-way phases

The Celestial Mechanics Approach (CMA)



Within the Bernese GNSS Software (BSW):

- Numerical integration (with a priori parameters) of equations of motion and variational equations.
- Set up of normal equations (NEQs) **for each observation type** (Doppler, KBRR, positions, ...) on a daily basis.
- Combination of the NEQs with appropriate weighting.
- NEQ manipulation (e.g., preelimination of parameters and accumulation to weekly, monthly, etc... NEQs)
- NEQ inversion \rightarrow simultaneous solution for the improved parameters (orbit, gravity field coefficients, etc...)




Orbit propagation

- Daily arcs, initial conditions from GNI1B science orbits
- Force model (SGM150J and GRGM900C up to d/o 600 a priori)




Parameter estimation

Observation	nObs	error	weight	
2-way Doppler	416746	0.3 mm/s	1	
1-way Doppler	334181	0.08 mm/s	0.1	
KBRR	1261440	0.03 $\mu\text{m/s}$	10^8	



Parameter		R	S	W
Arc-specific				
x_0, v_0	6			
emp. accel.	opr	×		
pulses	30'	0	1 $\mu\text{m/s}$	1 $\mu\text{m/s}$
Global				
Gravity field coeff.	×			
Love number k_2	×			

Orbit determination: KBRR residuals




Visible effects:

- impact of the pseudo-stochastic pulses (every 30');
• solar radiation pressure at light/shadow transitions.

Based on GRGM900C, d/o 600

Orbit determination: KBRR residuals





Visible effects:

- impact of the pseudo-stochastic pulses (every 30');
- solar radiation pressure at light/shadow transitions.


Huge improvement from additional SRP and PRP modeling!!

Orbit determination: improved eclipses


No SRP, ERP
(pulses + empirical only)



Orbit determination: improved eclipses




RMS [$\mu\text{m/s}$]




RMS [$\mu\text{m/s}$]

RMS [$\mu\text{m/s}$]





RMS [$\mu\text{m/s}$]

SRP + ERP




Orbit determination: improved eclipses

Solar Panel Current based




Orbit determination: improved eclipses

Solar Panel + Time Shift




0.0 0.1 0.2 0.3 0.4 0.5




0.0 0.1 0.2 0.3 0.4 0.5

RMS [$\mu\text{m/s}$]






0.0 0.1 0.2 0.3 0.4 0.5
RMS [$\mu\text{m/s}$]



Credits: NASA/JPL

Gravity field determination


- **A priori orbit: fit of Doppler and KBRR observations with appropriate parametrization**
- Combination with KBRR daily NEQs
- NEQs stacking up to the whole mission (pre-elimination of orbit parameters)
- NEQs inversion and solution



$$V(r, \lambda, \phi) = \frac{GM_M}{r} \sum_{l=1}^{l_{\max}} \left(\frac{R_M}{r} \right)^l \sum_{m=0}^l \bar{P}_{lm}(\sin \phi) (\bar{C}_{lm} \cos m\lambda + \bar{S}_{lm} \sin m\lambda)$$

Gravity field determination: GRGM900C


Difference degree amplitudes : $\Delta_l = \sqrt{\frac{1}{2l+1} \sum_{m=0}^l (\Delta \bar{C}_{lm}^2 + \Delta \bar{S}_{lm}^2)}$



a priori GRGM900C
(d/o 660): 2-way
Doppler + KBRR

Gravity field determination: GRGM900C


Difference degree amplitudes : $\Delta_l = \sqrt{\frac{1}{2l+1} \sum_{m=0}^l (\Delta \bar{C}_{lm}^2 + \Delta \bar{S}_{lm}^2)}$



a priori GRGM900C
(d/o 660): 1-way
Doppler + KBRR

Gravity field determination: GRGM900C

Difference degree amplitudes : $\Delta_l = \sqrt{\frac{1}{2l+1} \sum_{m=0}^l (\Delta \bar{C}_{lm}^2 + \Delta \bar{S}_{lm}^2)}$




NEQ Combination:

$0.1 \times 1\text{-way} +$
 $1 \times 2\text{-way} +$
 $10^8 \times \text{KBRR}$


Gravity field determination: GRGM900C

Difference degree amplitudes : $\Delta_l = \sqrt{\frac{1}{2l+1} \sum_{m=0}^l (\Delta \bar{C}_{lm}^2 + \Delta \bar{S}_{lm}^2)}$




Gravity field determination: GRGM900C

Gravity field anomalies AIUB-GRL350B





Solution	$k_2 \pm \sigma$
AIUB-GRL350B	0.024147 ± 0.000069
GRGM900C	0.024116 ± 0.000108


Gravity field determination: SGM150J





a priori SGM150J




Gravity field determination: SGM150J



a priori SGM150J
 1 iteration: April





Gravity field determination: SGM150J




a priori SGM150J

1 iteration: April

2 iteration: 62-140



Gravity field determination: SGM150J





a priori SGM150J

1 iteration: April


2 iteration: 62-140

3 iteration:

SRP+PRP modeling



Gravity field determination: SGM150J



a priori SGM150J

1 iteration: April



2 iteration: 62-140

3 iteration:


SRP+PRP modeling

4 iteration:

larger param. space



Gravity field determination: SGM150J



a priori SGM150J

1 iteration: April

2 iteration: 62-140



3 iteration:

SRP+PRP modeling

4 iteration:

5 iteration: 62-149

larger param. space




GRAIL + Lunar Laser Ranging

Combination of techniques \Rightarrow Optimal solutions for geophysics

- GRAIL (AIUB) + LLR (POLAC@SYRTE)

Collaboration with A. Bourgoin



Gravity field, tides, rotation




Low degrees gravity field, tides,
rotation, Earth/Moon distance

- Combination on NEQ level (full covariance information)
- Different geometries \Rightarrow different sensitivity
- Similar principle as GRACE + SLR on Earth
- **Current status:** combined gravity + k_2


Conclusions

- Bernese GNSS Software capable of 1-way and 2-way DSN Doppler data → **inhouse** processing of planetary mission data
- Parallelized MKL/Lapack environment for large solutions
- Final Moon gravity field (and tides) solution on track
- Collaborations and ongoing projects in planetary geodesy @ AIUB


Venus



Mercury



Europa





Credits: JAXA (Selene)

Thanks for your attention!
Questions?

LAPACK/INTEL MKL

- High resolution Moon gravity fields = high computational needs
- Parallel code for gravity NEQ setup (NEQSYS2) + inversion (NEQSOLVE)

d/o	n. of parameters	standard	MKL
200	40397	≈ 1 week	≈ 1 day (20' Y)
300	90527	≈ 2.5 week	≈ 3 days (1.5 h Y)
350	123000	-	≈ 1 week (3.5 h Y)



UBELIX


+



Parallel Intel MKL/LAPACK libraries for optimal use of parallel
UniBE HPC Cluster (UBELIX) environment

Combined orbit: KBRR residuals




0.68 mm/s



Daily RMS values of KBRR residuals for GRAIL-A (PM), using different gravity field models and parametrizations.

- Days 140 – 150 at lower altitude → larger residuals

Observed vs computed : Doppler residuals



34 mHz

7 mHz

2 mHz

Non-gravitational forces: setup

- Solar Radiation Pressure (28-plates macromodel, Fahnestock 2012)






Figure 1. Placement of surfaces on GRAIL-B in highest fidelity model.

Non-gravitational forces: setup

- Solar Radiation Pressure (28-plates macromodel, Fahnestock 2012)
- Moon Albedo (Delft Albedo Model-1, d/o 15)



- Moon IR (black body)

